



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES—UNTABULATED.

BRAZIL—Ceara.—Month of October, 1909. Estimated population, 50,000. Total number of deaths 81, including tuberculosis 12, typhoid fever 7.

COSTA RICA—San Jose.—Month of November, 1909. Estimated population, 27,532. Total number of deaths 52, including diphtheria 1, tuberculosis 2.

CUBA—Matanzas.—Month of November, 1909. Estimated population, 36,009. Total number of deaths 50, including diphtheria 1, tuberculosis 8.

CURAÇAO.—Five weeks ended December 3, 1909. Estimated population, 30,000. Total number of deaths, 33. No contagious diseases.

FRANCE—Calais.—Month of December, 1909. Estimated population, 80,000. Total number of deaths 107, including diphtheria 2, scarlet fever 1, tuberculosis 30, typhoid fever 1.

Cherbourg.—Month of November, 1909. Estimated population, 43,837. Total number of deaths, 38, including tuberculosis 24, typhoid fever 4.

Nice.—Month of October, 1909. Estimated population, 150,881. Total number of deaths, 168, including tuberculosis 26, typhoid fever 6.

Roubaix.—Month of November, 1909. Estimated population, 121,115. Total number of deaths, 153, including diphtheria 2, tuberculosis 36, typhoid fever 1.

St. Etienne.—Two weeks ended November 30, 1909. Estimated population, 150,000. Total number of deaths, 137, including tuberculosis 19, typhoid fever 1.

GREAT BRITAIN AND IRELAND.—Week ended November 27.

England and Wales.—The deaths registered in 76 great towns having an aggregate estimated population of 16,445,281 correspond to an annual rate of 16.4 per 1,000.

Ireland.—The deaths registered in 21 principal town districts having an aggregate estimated population of 1,442,830 correspond to an average annual rate of 21 per 1,000. The lowest rate was recorded in Tralee, viz, 5.3, and the highest in Newtownards, viz, 28.6.

Scotland.—The deaths registered in 8 principal towns having an aggregate estimated population of 1,839,803 correspond to an annual rate of 24.2 per 1,000. The total number of deaths recorded was 866, including diphtheria 13, measles 49, scarlet fever 7, typhoid fever 3, whooping cough 9. The lowest rate was recorded in Greenock, viz, 11.5, and the highest in Glasgow, viz, 32.7.

MAURITIUS.—Month of September, 1909. Census population, 374,237. Total number of deaths, 1,279, including plague 38, tuberculosis 81, typhoid fever 4.